
# **Zuclopentixol Injection**

# This is an easy way to read this word... zoo-clo-pen-tick-sol



#### What is this leaflet for?

This leaflet is to help you understand more about your medicine.



### What is the medicine called?

Zuclopentixol is also called Clopixol.

You will see 1 of these names on your prescription.

Zuclopentixol will come as an injection.



## What is zuclopentixol for?

Zuclopenhixol could be used for different reasons.

Zuclopentixol is used for psychoses and mania.

Zuclopentixol is sometimes used for people who have behaviour problems.

For example, it could...



- help if you are anxious
- help calm you down
- help you feel less angry
- help you feel less tense.



## What will zuclopentixol do?

Zuclopentixol should help you feel calm and relaxed.

It could take some time for zuclopentixol to have its full effect.

This effect should reduce your behaviour problem.



For example, it may...

- help you control your temper
- help you stop hurting yourself
- help you stop breaking things.



## What are the side effects of zuclopentixol?

Side effects are effects medicine can have on your body that are not wanted.

Not everyone taking a medicine will get side effects.

Many side effects will go away with time.

Many side effects are rare.



You might want to talk to your doctor if you have any side effects.

For example, these are some side effects of zuclopentixol...

- feeling sleepy or sluggish
- feeling more restless or agitated
- dry mouth
- too much spit
- headache
- you could lose interest in sex
- putting on weight
- constipation
- blurred vision or things look fuzzy
- if you are female your periods could change.







Some side effects are more serious.

You should tell someone straight away if...

- you feel faint or dizzy
- you have a fever or high body temperature
- you feel stiff or shaky
- you have problems with weeing.



How and when do I take zuclopentixol?

A doctor or nurse will give you your injection.



The injection will usually be given in your buttock.

You will normally be given an injection every 2 weeks.



If you forget to have your injection, tell someone straight away.

You should carry on having your injection even if you feel better unless your doctor asks you to stop.

You should talk to your doctor before you stop taking your medicine.



#### What do I need to know about this medicine?

Drinking alcohol whilst taking zuclopentixol could make you feel drowsy.

Some medicines do not work well together.

You should tell your doctor about any other medicines you are taking.



## Where can I get more information?

This leaflet does not tell you everything about zuclopentixol.

If you want more information about zuclopentixol...

- you can ask your pharmacist, doctor or nurse for more information
- you can ask your carer to get more information
- you can phone NHS Direct on 0845 4647
- you can phone a medicine information helpline on 020 3228 2999.

G. Unwin and S. Deb from the University of Birmingham created this leaflet with help from People First (Staffs) and Mencap. The project was funded by the Big Lottery Fund.

It can be downloaded from www.ld-medication.bham.ac.uk.

The information in this leaflet came from...

- the British National Formulary (www.bnf.org)
- the United Kingdom Psychiatric Pharmacy Group Patient Advisory Leaflets (www.ukppg.org.uk/ukppg-pals.html)
- Miriam Wilcher's Handbook for Carers of People with Learning Disabilities.